

AD _____

GRANT NUMBER DAMD17-96-1-6238

TITLE: Regulation of Growth Factor Release By Proteases
Inhibitors in Breast Cancer

PRINCIPAL INVESTIGATOR: Thomas H. Finlay, Ph.D.

CONTRACTING ORGANIZATION: New York University Medical Center
New York, New York 10016

REPORT DATE: September 1997

19980116 160

TYPE OF REPORT: Annual

PREPARED FOR: Commander
U.S. Army Medical Research and Materiel Command
Fort Detrick, Frederick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for public release;
distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

DTIC QUALITY INSPECTED 3

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE September 1997	3. REPORT TYPE AND DATES COVERED Annual (15 Aug 96 - 14 Aug 97)	
4. TITLE AND SUBTITLE Regulation of Growth Factor Release By Proteases Inhibitors in Breast Cancer			5. FUNDING NUMBERS DAMD17-96-1-6238	
6. AUTHOR(S) Thomas H. Finlay, Ph.D.				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) New York University Medical Center New York, New York 10016			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Commander U.S. Army Medical Research and Materiel Command Fort Detrick, Frederick, Maryland 21702-5012			10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200) This report describes the cloning, sequencing and properties of a potential TGF α -releasing protease present in human breast epithelial cells. Degenerate oligonucleotides based on the conserved sequences about the his ⁵⁷ and ser ¹⁹⁵ residues in mammalian serine proteases were used to clone a novel serine protease sequence (PCR-7) from an MCF-7 breast cancer cell cDNA library. The deduced 321 amino acid open reading frame shows a high degree of homology with the cell-surface serine proteases enterokinase, prekalikrein, hepsin and prostasin. Northern blot analysis, using a PCR-7 cDNA probe, shows the presence of an approximately 4.2 kb mRNA species in total RNA from MCF-7 cells. Similar mRNA species were found in normal breast tissue, trophoblast, and uterine glandular epithelium but not in human U373MG glioma or HepG2 hepatoma cells. In MCF-7 cells, expression of the PCR-7 mRNA was not significantly stimulated by estradiol or TGF α , but was stimulated by IL-6 (200%) and IL-1 (300%). A GST fusion protein constructed from a PCR-7 cDNA sequence was used to raise antibodies in rabbits. On western blot analysis, the antibody recognized a 33 kDa polypeptide in trophoblast and MCF-7 cells under both reducing and non-reducing conditions. This antibody inhibited TPA-induced TGF α release from MCF-7 cells in a dose-dependent manner as compared to pre-immune sera.				
14. SUBJECT TERMS Breast Cancer MCF-7 Cells, alpha-1-Antitrypsin TGF-alpha, serine protease, cloning			15. NUMBER OF PAGES 18	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

 Where copyrighted material is quoted, permission has been obtained to use such material.

 Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

✓ Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

✓ In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Resources, National Research Council (NIH Publication No. 86-23, Revised 1985).

✓ For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

✓ In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

✓ In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

✓ In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.



PI - Signature

9/12/95

Date

Table of Contents

Front Cover

SF 298 Report Documentation Page

Foreword

Table of Contents

Introduction	5
Materials and Methods	6
Results/Discussion	9
Conclusions	16
References	17

CLONING OF A POTENTIAL TGF α -RELEASING SERINE PROTEASE FROM MCF-7 HUMAN BREAST CANCER CELLS

Susan S. Kadner and Thomas H. Finlay

Division of Reproductive Biology, Department of Obstetrics and Gynecology
and the Kaplan Cancer Center
NYU Medical Center, New York, NY 10016

Supported by grant DAMD-17-96-1-6238 from the
US Army Medical Research and Material Command

INTRODUCTION

The broad aim of this proposal is to test the novel hypothesis that the coordinate action of proteases and protease inhibitors are responsible for growth factor activation and release in the human breast cancer cell, and for the progression of breast cancer *in vivo*. One-third of all cases of advanced breast cancer are estrogen responsive, and recent epidemiological studies and studies using human breast cancer cells in culture strongly suggest a correlation between estrogens and the pathogenesis of breast cancer (Harris et al 1992). The mechanisms by which this occurs are not entirely clear. Certainly, estrogens have a direct effect on cell growth. In addition, they can stimulate the expression and release of a variety of polypeptide growth factors. It is highly likely that the tumorigenic effects of estrogens are due, at least in part, to the autocrine/paracrine action of these factors. Several of these polypeptides, including epidermal growth factor (EGF), and its analogs heregulin and transforming growth factor- α (TGF α) and the insulin-like growth factors (IGF-I and IGF-II), have been shown to require pericellular proteolysis for activation or release (Massagué and Pandiella 1993). To achieve homeostasis in a normal breast epithelial cell, levels of these pericellular growth factor activating proteases also must be regulated. We hypothesized that this was accomplished by the action of locally synthesized protease inhibitors. Thus an imbalance in the ratio between local levels of particular proteases and protease inhibitors could be responsible for increases in tumorigenic potential.

TGF α , a peptide structurally and functionally related to EGF, interacts with the EGF receptor and elicits a mitogenic response in a variety of cells (Lee et al 1995). TGF α expression occurs in normal breast tissue, breast tumors and breast cancer cells in culture and TGF α has been proposed to act as a major autocrine mediator of estrogen-stimulated growth in estrogen-dependent breast cancer cells (Harris et al 1992). TGF α is synthesized as part of a glycosylated transmembrane precursor (proTGF α) which can be processed intracellularly and extracellularly by glycosylation and proteolysis to yield a family of polypeptides ranging in size from the 29kDa transmembrane proTGF α to the 6 kDa soluble, mature form (Massagué and Pandiella 1993). Complete processing does not occur to the same extent in all tissues. While all

TGF α forms appear to possess some degree of biological activity, there is good evidence that particular biological actions may depend on the degree of proteolytic processing. The proteolytic cleavage required for release of soluble TGF α from its membrane-bound precursor on the cell surface may be an essential step in regulating TGF α activity.

Results from this laboratory have clearly demonstrated a relationship between tumorigenicity of MCF-7 human breast cancer cells (as measured by growth in soft agar), endogenous synthesis of the protease inhibitor α_1 -antitrypsin (α_1 -AT) and release of TGF α (Tamir et al 1990, Finlay et al 1993a,b). Growth in soft agar was blocked by α_1 -AT whether added to the tissue culture media or synthesized by the tumor cell itself. A useful tool in these studies was a new MCF-7 cell subline, producing 10-fold higher levels of α_1 -AT than its parental cell line, constructed by stable transfection with an α_1 -AT cDNA (Yavelow et al 1997). Growth in soft agar and release of TGF α was decreased in cells transfected with the α_1 -AT cDNA when compared to cells transfected with vector alone. Consistent with the above we had identified a serine protease with elastase-like activity, capable of forming a stable complex with α_1 -AT, on the MCF-7 cell surface. In the work described here, we have focused our attention on the identification of the pericellular protease responsible for TGF α release, as this might be a convenient target for anticancer drug action.

METHODS AND MATERIALS

Preparation of cDNA - Total cellular RNA was prepared from MCF-7 (ML) cells by extraction in guanidinium isothiocyanate and centrifugation on cesium chloride. mRNA was isolated using PolyATtract mRNA isolation system IV (Promega) and used to synthesize double stranded cDNA (BRL cDNA synthesis system).

Primers - Degenerate oligonucleotide primers designed from the conserved amino acids around the active site residues his⁵⁷ and ser¹⁹⁵ and the oligonucleotide probe based on the conserved sequence around asp¹⁰² were synthesized as previously described (Sakanari et al, 1989; Elvin et al, 1993).

PCR conditions - 100 ng of cDNA was amplified in a Perkin Elmer thermal cycler 480 as described by Elvin, et al except that 4 mM MgCl₂ was used in the reaction.

Cloning, Southern blotting and sequencing of PCR products - 2 μ l of the PCR products were ligated directly into the TA cloning vector pCRTMII (Invitrogen). 1 μ l of the ligation reaction was used to transform E. Coli strain INV α F' (Invitrogen). Recombinant clones were selected by blue / white color selection on LB agar plates containing ampicillin (50 μ g/ml) and X-gal. Plasmid DNA was isolated from 22 white colonies by alkaline lysis, digested with Eco RI and electrophoresed on a 1.2% agarose gel containing 0.5 μ g/ml ethidium bromide. After denaturation and neutralization the gel was transferred to

GeneScreen nylon membrane (NEN), baked at 80° C for two hours and hybridized overnight at 37° C with the degenerate Asp¹⁰² probe end labeled with γ -³²P ATP using T4 polynucleotide kinase (Promega). The blot was washed with 4 X SSC, 0.1% SDS for 2 X 5 min at room temperature followed by 40 min at 37° and briefly at room temperature. One strongly hybridizing fragment of \approx 460 bp, (PCR-7), was gel purified and used as a probe for northern blotting. Plasmids were sequenced using SP6 and T7 promoter primers and Sequenase DNA sequencing kit (USB).

Northern Blots - Total RNA, isolated from MCF-7 cells, U373MG glioma cells, Hep G2 cells, normal human breast tissue, trophoblast and uterine glands was electrophoresed on 1.5% agarose/formaldehyde gels, transferred to GeneScreen and hybridized with ³²P-labeled probe. All probes used for northern blots were labeled by Megaprime DNA labeling system (Amersham).

Preparation of cDNA library - Total RNA was extracted from MCF-7 cells using Ultraspec RNA isolation system (Biotecx). mRNA was isolated using the MGP mRNA Purification Kit (CPG). Starting with 3.8 μ g of mRNA, a cDNA library was constructed in the plasmid vector pSport 1 using the SuperScript Plasmid System for cDNA Synthesis and Plasmid Cloning (Life Technologies) according to the manufacturers instructions. The library was constructed directionally in the Not I-Sal I cut vector such that cloned genes can be expressed *in vitro* using the T7 RNA polymerase promoter. After transformation of E. Coli DH5 α the library was amplified on selective agar plates, aliquotted and stored at -70° C.

Screening of cDNA library. Approximately 60,000 recombinant colonies were screened by standard hybridization techniques after transfer to Nytran nylon membranes (Schleicher & Schuell). In order to eliminate any potential background hybridization to plasmid sequences, the 460bp protease insert was gel purified and used as a template for PCR. The primers, determined from the sequence data, amplified a 300bp fragment which was purified and used to probe the library. This probe does not hybridize to plasmid DNA as determined by Southern blot. Positive colonies were selected and rescreened. Insert size was determined by electrophoresis of Not I - Sal I cut plasmid DNA.

Preparation of Fusion Protein and Antibody Production. The 460bp PCR-7 protease fragment was ligated to the Eco RI site of the glutathione S-transferase gene fusion vector pGEX-4T-1 (Pharmacia). The resulting construct as well as the parent vector were used to transform E. Coli strain INV α F', a strain recommended for cloning and maintenance of the plasmids. Selected colonies were grown in liquid culture for 2.5 h at 37° then induced with 0.1mM isopropyl- β -D-thiogalactoside (IPTG) for an additional hour. Cell pellets were suspended in ice-cold PBS, sonicated briefly and electrophoresed on a 12% SDS polyacrylamide gel. After determining which clone was in the right orientation for expression of the GST-PCR-7 fusion protein, the plasmid (pGEX-4T-1-PCR-7) was used to transform E. Coli strain BL21 for large-scale

expression. Fusion protein was purified from 250 ml of an induced culture by the procedure described by Frangioni (1993) for insoluble fusion proteins. Polyclonal antibodies were produced in rabbits by the Pocono Rabbit Farm & Laboratory, Inc. IgG was purified by ammonium sulfate precipitation followed by ion-exchange chromatography on DEAE-cellulose. Western blots were visualized by autoradiography after detection with ^{125}I protein A.

RESULTS AND DISCUSSION

Taking advantage of the high degree of homology in the DNA sequences about the active site histidine, aspartate and serine residues (his⁵⁷, asp¹⁰², ser¹⁹⁵) in all serine proteases, we constructed degenerate oligonucleotide primers which were used to "fish out" putative serine protease sequences from an MCF-7 cell cDNA library. In the initial step, a series of approximately 500 bp fragments between the his⁵⁷ and ser¹⁹⁵ sites were amplified by PCR. The amplified sequences were then cloned into a PCR cloning vector, which was used to construct a mini-cDNA library. Clones from the mini library were selected by Southern blotting using a ³²P-labeled degenerate oligonucleotide probe based on the sequence about asp¹⁰². Several positive clones (Fig. 1) were sequenced. The 460 bp sequence of one strongly hybridizing clone, PCR-7, showed a high degree of homology to known serine proteases (64% identity in a 220 bp region at the 3' terminus to human trypsinogen-B and a 60% identity in a 120 bp region around the 5' terminus to human pancreatic protease).




Figure 1. Cloning of MCF-7 Serine Protease Fragments. Putative cDNA inserts corresponding to the his⁵⁷-ser¹⁹⁵ sequence were excised from the pcR11 cloning vectors, electrophoresed on agarose gels and stained with ethidium bromide (panel A). The DNA fragments were then transferred to nitrocellulose and probed with a ³²P-labeled oligonucleotide corresponding to the sequence about asp¹⁰² (panel B). While all of the clones contained inserts between 400 and 600 bp, in this series only the clone-7 insert (4th lane from right) showed significant hybridization to the asp¹⁰² probe.

Northern blot analysis, using a PCR-7 cDNA probe, showed the presence of an approximately 4.2 kb mRNA species in total RNA from MCF-7 cells. This probe also recognized similar mRNA species in normal breast tissue, trophoblast, and proliferative phase uterine glandular epithelium but not in human U373MG glioma or HepG2 hepatoma cells (Fig. 2). In trophoblast isolated from term placenta, expression of the PCR-7 mRNA was repressed when trophoblast was maintained under hypoxia for 48 h. We have shown that under these conditions, TGF α release is reduced. In MCF-7 cells, expression of the PCR-7 mRNA was not significantly stimulated by estradiol or TGF α , but was stimulated by TGF β (150%), IL-6 (200%) and IL-1 (300%) (Fig. 3).




Figure 2. Northern Blot Analysis Using Clone-7 cDNA Probe. Total RNA was prepared from various human tissues and cell lines, electrophoresed on a 1.5% agarose/formaldehyde gel, transferred to GeneScreen nylon membrane and probed with the ^{32}P -labelled cDNA insert. The RNAs shown are lane 1, human breast ductal epithelium; lane 2, MCF-7 ML cells; lane 3, MCF-7 P203 cells; lane 4, U373MG glioma cells; lane 5, term trophoblast cultured under hypoxic conditions; lane 6, term trophoblast cultured under normoxic conditions; lane 7, MCF-7 cells in media containing 10^{-8} M estradiol, lane 8, MCF-7 cells in estradiol-free media; lane 9, uterine glandular epithelium.




Figure 3. Northern Blot Analysis of MCF-7 Cell RNA. Effect of Growth Factors and Cytokines on PCR-7 mRNA Expression.

MCF-7 cells were maintained 48 hours in phenol red-free DMEM containing 10% stripped FBS. The media was replaced with fresh media containing the indicated additions and incubated 24 hours longer. Total RNA (ca. 15 μg) from each sample was subjected to Northern blot analysis with a ^{32}P -labeled PCR-7 cDNA. The blot was reprobed with an actin cDNA. lane 1, no additions; lane 2, 10^{-8} M Estradiol. lane 3, 50 ng/ml TGF- α ; lane 4, 5 ng/ml TGF- β ; lane 5, 10 u/ml IL-1; lane 6, 200 u/ml IL-6. The blot was reprobed with a ^{32}P -labeled actin cDNA (insert). The blot was densitometrically scanned and integrated

The 460 bp PCR-7 cDNA was also ligated into the glutathione S-transferase (GST) gene fusion vector, pGEX-4T-1. This construct and the parental vector were used to transform *E. coli* and fusion protein expression was induced with IPTG. On SDS PAGE, 3 of 6 recombinant vectors displayed a prominent band of about 40 kDa representing the GST- PCR-7 fusion protein while the parental vector produced a prominent 29 kDa band characteristic of GST alone. Large scale cultures of bacteria containing the GST- PCR-7 fusion protein vector were prepared and substantial amounts of the GST- PCR-7 fusion protein was isolated. The fusion protein was used to raise antibodies in two rabbits. Antiserum from one of the rabbits was able to

immunoprecipitate the ^{35}S -labeled proteins prepared in vitro described below. Preliminary evidence suggests that this antibody is able to inhibit TPA-induced $\text{TGF}\alpha$ release from MCF-7 cells in a dose-dependent manner as compared to pre-immune sera (Fig. 4).




Figure 4. Antibody to PCR-7 fusion protein blocks release of $\text{TGF}\alpha$ from MCF-7 cells.

MCF-7 (ML) cells were maintained in phenol red - free DMEM containing 10% FBS.

Estradiol (10^{-8} M) was added for 24 hours. Cells were then washed twice with serum-free media and allowed to condition serum-free DMEM containing estradiol and IgG for 30 min. TPA (50 ng/ml) was added and the cells were incubated for an additional 4 hours. Conditioned media were concentrated on C_{18} SEP-columns equilibrated with 10% acetonitrile-0.1% TFA eluting with 40% acetonitrile-0.1% TFA. Samples were taken to dryness and resuspended in PBS. $\text{TGF}\alpha$ levels were determined using a $\text{TGF}\alpha$ ELISA assay (Oncogene Science) according to the manufacturer's instructions. Results are reported as the percentage of $\text{TGF}\alpha$ released in the presence of antibody (IgG) to the PCR-7-GST fusion protein as compared to the same level of pre-immune IgG.

Two cDNAs of approximately 3.8 and 3.2 kb containing the PCR-7 sequence were cloned from a second MCF-7 cell cDNA library using the PCR-7 sequence as a probe. The library had been constructed by directional cloning into the multiple cloning site of a pSport I vector such that the 5' terminus was downstream from a T7 promoter sequence. Except for minor differences in the 3' sequence, the two clones appear to be identical where they overlap. Both clones hybridized to the same size mRNA as did the PCR-7 sequence used to screen the library (Fig. 5). Approximately 2000 bp (plus an additional 500 bp at the 5' end) of the larger clone (Clone 1) has been sequenced (Fig. 6). The sequence has a 963 b open reading frame (bases 463-1426) and an approximately 500 b 3' sequence containing the poly adenylation site. The deduced amino acid sequence from the open reading frame contains the His⁵⁷ and Ser¹⁹⁵ motifs characteristic of serine proteases (amino acid residues 119-124 and 265-275 respectively). The aspartate residue corresponding to Asp¹⁰², the third amino acid in the serine protease catalytic triad, is most likely one of the three Asp residues between residues 171 and 177. With BLAST, a high degree of identity at the amino acid level was found between the PCR-7 protease sequence and the serine proteases (or their zymogens) enterokinase precursor (Kitamoto et al 1995), hepsin (Leytus et al 1988), prekalikrein (Chung et al 1986), coagulation factor XI (Fujikawa et al 1986), acrosin precursor (Baba et al 1989) and prostasin (Yu et al 1994, 1995) (Table I). Significantly, enterokinase precursor, hepsin, prekalikrein, acrosin precursor and prostasin are cell surface bound. A high degree of homology was found at the nucleotide level with another cell surface protease, hepatocyte growth factor activator (Miyazawa et al 1993).




Figure 5. Full Length cDNAs (Clone 1, Clone 2) and 300 bp PCR Fragment Used for Cloning Bind to the Same Size mRNA. The 3.8 (Clone 1) and 3.2 (Clone 2) kb cDNAs containing the PCR-7 sequence were gel purified, labeled with ^{32}P -dCTP and used to probe a northern blot containing 15 μg of ML total RNA / lane. The 300 bp PCR-7 cDNA probe used to screen the library was used as a control.

Table 1. Sequence comparison analysis of PCR-7 Protease with other human serine proteases. Analysis was performed at the NCBI using the BLAST network service.

Protein	% Identity	Overlap
Enterokinase Precursor	40.6	254 aa
Hepsin	39.6	270 aa
Prekallikrein	38.8	260 aa
Factor XI	35.8	327 aa
Acrosin Precursor	35.5	276 aa
Prostasin Precursor	51.9	187 aa
Hepatocyte Growth Factor Activator	53.5	620 bp (740-1360)

Figure 6. Sequence of PCR-7 Protease cDNA. Clones were sequenced at the NYU/Skirball sequencing facility using an Applied Biosystems 373 automated DNA sequencer. Sequencing primers were obtained from Gibco BRL. Deduced sequence of the 321 amino acid open reading frame (bases 463-1426) is shown beneath the corresponding trinucleotide.

1	gcgcttcaaatcttcttaacotgctggagccggcggtgctgcgggcaactgccccaaag		
61	antacgtggagatcaacgggggagaataactgcggagagaggtccagttctgctgtcacca		
121	gcaacagacaacaagatcacagttcgcttccactcagatcagtcctacacggacacggct		
181	tattgacacctctagttgccccctctttagacgacctctctccaggttcaagcagcagtggt		
241	gcaacagacaacaagatcacagttcgcttccactcagatcagtcctacacggacacggct		
301	tattgacacctctagttgccccctctttagacgacctctctccaggttcaagcagcagtggt		
361	gcaacagacaacaagatcacagttcgcttccactcagatcagtcctacacggacacggct		
421	tattgacacctctagttgccccctctttagacgacctctctccaggttcaagcagcagtggt		
481	AAAGCCAGCAGTGAATGGGAAGGACGACTGTGGGACGGTTCCGACGAGGCTCCTGC		
541	CCCAAGGTGAACGTGCTCACTTGTACCAACACACATACCCTGCCTCAATGGGCTCTGC		
601	TTAGAGCAAGGGCAACCTGAGTGTGACGGGAAGGAGGACTGTAGCAGCGCTCAGATGAG		
661	AAGGACTGCGACTGTTGGCTGCGGTCATTCACGAGACAGGCTCGTGTGTTGGGGGCACG		
721	GATGCGGATGAGGGCGAGTGGCCCTGGCAGGTAAAGCTGCATGCTTGGGCCAGGGCCAC		
781	ATCTCGGTGCTTCCTCATCTCTCCAACTGGTGGTCTCTGCGGCACACTGCTACATC		
841	GATGACAGAGGATTCAAGTACTCAGACCCACGACGagggccttctTGGGCTTGACGCA		
901	CCAGAGCCAGCGCAGCGCCCTGGGTCGAGGAGCGCAGGCTCAAGCGCATCATCTCCAC		
961	GGTCTCGGTGCGCTCGGGGACCCACGTCCTCGCGTCCGAGTTCGCGTAGTAGAGGGT		
1021	CGTCTCATGTGCGAGGTACCACGCGGGTAGACGACGGCTCGGAGGGTACAGAAGGA		
1081	GCCGGCAAGGCCATCTGGGTACGGGCTGGGGACACACCCAGTATGAGGACCTGGCGG		
1141	CTGATCCTGCAAAAGGGTGAGATCCGCGTCATCAACAGACACCTGCGAGAACCTCTCTG		
1201	GGCGTCGTTTGTAGTGGCGCGCTACTACACGACCCGAAGGAGTCCGCGCCGACCTGAGG		
1261	TGCCAGGGTGATTCCGGGGACCCCTGTCCAGCGTGGAGGCGGATGGGCGGATCTCCAG		
1321	GCCGGTGTGGTGAGCTGGGGAGACGGCTGCGCTCAGAGGAACAAGCCAGGCGGTACACA		
1381	AGGCTCCCTCTGTTTCGGGACTGGATCAAGAGAACACTGGGGTAtagggggcggggcca		
1441	cccaaatgtgtacacctgcgggggccaccatgctccacccagtggtgacgctgcaggc		
1501	tgagagctggacgctgactgacccagcgcccccagaacatacactgtgaactcaatctc		
1561	aggtagaaggggttctactgaccccaactgggggcaaggttgaagacacagcttcccccg		
1621	ccagcccaagctggggcgaggcgctgtgtgtatctgctccctgctgtgaaggag		
1681	gggtgggggtcgacccggctccgagcaacacataagacggaggggagacagattctctc		
1741	gtgccccgtgctcgaagcctggaggagtgacttccaccccccagcgcttagaccog		
1801	acaccccggaacccgggtgcgagaactccttcgggttcgagcctcctgggacctttgtc		
1861	gtgtaaatgagtaaaacattttatttttttttttttttttttttttttttttttttttt		
1921	cacatttactcattttgtataaaataaangaaaaatttttttttttttttttttttttt		
	Trypsin His57 Motif: ValSerAlaAlaHisCys	119	124
	Trypsin Ser195 Motif: AspSerCysGlnGlyAspSerGlyGlyProLeu	265	275
	Asn Glycosylation Motif: AsnGlnThrThr	241	244




Figure 7. In vitro Transcription/Translation and Immunoprecipitation of Putative PCR-7 Protease Clones.

Plasmids containing the 3.8 kb (Clone 1) and 3.2 kb (Clone 2) inserts were translated in the presence of ^{35}S methionine using a coupled reticulocyte lysate transcription/translation system (Promega TNT). Aliquots of the translation products were immunoprecipitated with rabbit polyclonal antisera to the GST-PCR-7 fusion protein and electrophoresed on a 10% SDS polyacrylamide gel. 'Control' indicates translation mix less template.

Using a coupled reticulocyte lysate transcription/translation system in the presence of ^{35}S methionine, both clones expressed proteins of approximately 24, 33 and 70 kDa while the 3.8 kb clone expressed a fourth protein of about 80 kDa (Fig. 7). The nature of the 70 and 80 kDa bands is unclear. All were also immunoprecipitated by the antiserum raised against the fusion protein. However, in both MCF-7 cells and trophoblast, the major protein band seen on western blot analysis using this same antibody had an apparent MW of approximately 33 kDa under both reducing and non-reducing conditions (Fig. 8).




Figure 8. Western Blot Analysis of PCR-7 Protease in MCF-7 Cells and Trophoblast.

Monolayers of MCF-7 cells were harvested by scraping, washed with PBS and lysed in SDS gel loading buffer in the presence or absence of β -mercaptoethanol (as indicated). Trophoblasts were isolated from term placenta and cultured for 48 hours before harvesting and lysis under reducing conditions. Aliquots were electrophoresed on a 12% SDS polyacrylamide gel, transferred to nitrocellulose and incubated with a 1-200 dilution of rabbit antibody to the GST-PCR7 fusion protein. Purified GST-PCR7 fusion protein was used as a control. Immune complexes were detected with ^{125}I -Protein A.

MCF-7 cells were surface labeled with [^3H]-diisopropylfluorophosphate (DIFP) and immunoprecipitated with antibody to the GST-PCR-7 fusion protein. When total proteins were analyzed by SDS-polyacrylamide gel electrophoresis two labeled bands of approximately 32 and 70 kDa respectively were detected (Fig. 9). Pretreatment of the cells with O-tetradecanoyl-phorbol-13-acetate (TPA), which has previously been shown to stimulate the release of $\text{TGF}\alpha$ in MCF-7 cells in the presence of estradiol (Yavelow et al, 1997), had no apparent effect on the intensity of either band. Our failure to immunoprecipitate the ^3H -DIFP labeled PCR-7 protease in this preliminary experiment is disturbing as it might indicate that the protein is not on the cell surface. However, it is likely that the labeled bands represent mixtures of several cell surface serine proteases of similar sizes, of which the PCR-7 protease is only a very small fraction. Similarly, this same level of antibody failed to significantly inhibit release of $\text{TGF}\alpha$ (Fig. 4). We suspect that not enough specific antibody was present in either of these experiments. Both experiments will be repeated with higher levels of affinity purified antibody to the GST-PCR-7 fusion protein. This will be accomplished by affinity chromatography on immobilized fusion protein. After elution of the antibody from the matrix with the appropriate chaotropic buffer, specific antibody will be further purified by affinity chromatography on immobilized GST.




Figure 9. MCF-7 Cells Contain a Cell Surface Protease Which Reacts With ^3H DIFP. MCF-7 cells (two ca. 75% confluent T-25 flasks per experiment) were washed with serum-free medium and incubated for 30 min in the absence (lane 1) or presence (lane 2) of 50 ng/ml TPA. ^3H DIFP (12.5 $\mu\text{Ci}/\text{ml}$, 8.4

$\mu\text{Ci}/\text{mmol}$) was added to each flask and the incubation was allowed to continue an additional 60 min. Cells were washed with PBS, lysed in buffer A by passing repeatedly through a 27 1/2 gauge needle and spun for 5 min. in a microfuge at 4° to remove cell debris. After preclearing the cell lysates with pre-immune rabbit IgG and protein A agarose, the supernatants were incubated with 5 μl of specific IgG (24.5 $\mu\text{g}/\text{ml}$) for 2 hours. Immune complexes were collected with protein A agarose, washed 6 times with buffer A, heated for 5 min at 95°C and electrophoresed on a 12% SDS polyacrylamide gel. Aliquots of cell lysates before and after immunoprecipitation were also electrophoresed. After treatment with en 3 hance, the gel was dried and exposed to x-ray film for 15 days. Lanes 1 and 2, immunoprecipitates; lanes 3 and 4, lysates after immunoprecipitation; lanes 5 and 6, lysates before immunoprecipitation. Lanes 2, 4, and 6 represent samples treated with TPA.

CONCLUSIONS

1. Degenerate oligonucleotides based on the conserved sequences about the his⁵⁷ and ser¹⁹⁵ residues in mammalian serine proteases were used to clone a potential TGF α -releasing serine protease (PCR-7) from an MCF-7 breast cancer cell cDNA library.
2. Northern blot analysis, using a PCR-7 cDNA probe, shows the presence of an approximately 4.2 kb mRNA species in total RNA from MCF-7 cells. Similar mRNA species were found in normal breast tissue, trophoblast, and proliferative phase uterine glandular epithelium but not in human U373MG glioma or HepG2 hepatoma cells. In MCF-7 cells, expression of the PCR-7 mRNA was not significantly stimulated by estradiol or TGF α , but was stimulated by IL-6 (200%) and IL-1 (300%).
3. A GST fusion protein constructed from a PCR-7 cDNA sequence was used to raise antibodies in rabbits. On western blot analysis, the antibody recognized a 33 kDa polypeptide in trophoblast and MCF-7 cells under both reducing and non-reducing conditions. This antibody inhibited TPA-induced TGF α release from MCF-7 cells in a dose-dependent manner as compared to pre-immune sera.
4. Because of the high degree of sequence homology with known membrane-bound serine proteases particularly, enterokinase, kininogen, hepsin and prostatic, and because antibodies to a PCR-7-GST fusion protein inhibit TGF α , we believe the PCR-7 sequence to be a membrane-bound TGF α releasing enzyme.

REFERENCES

- Baba, T., Watanabe, K., Kashiwabara, S-I and Arai, Y. 1989 Primary structure of human proacrosin deduced from its cDNA sequence FEBS Lett. 244:296-300.
- Chung, D.W., Fujikawa, K., McMullen, B.A. and Davie, E.W. (1986) Human plasma prekallikrein, a zymogen to a serine protease that contains four tandem repeats. Biochemistry 25: 2410-2417.
- Elvin, C.M., Whan, V. and Riddles, P.W. 1993 A Family of Serine Protease Genes Expressed in Adult Buffalo Fly. (*Haematobia irritans exigua*) Mol. Gen. Genet. 240:132-139.
- Finlay, T.H., Tamir, S., Kadner, S.S., Yavelow, J. and Levitz, M. 1993a α_1 -Antitrypsin and Anchorage-Independent Growth of MCF-7 Breast Cancer Cells. Endocrinology 133:996-1002.
- Finlay, T.H., Kadner, S.S. and Tamir, S. 1993b Protease Inhibitor Synthesis by MCF-7 Breast Cancer Cells. in Protease Inhibitors as Cancer Chemopreventative Agents. (W. Troll and A. Kennedy, Eds.). Plenum Press, New York. pp 141-159.
- Frangioni, J.V. and Neel, B.G. (1993) Solubilization and Purification of Enzymatically Active Glutathione S-Transferase (pGEX) Fusion Proteins. Analyt Biochem 210:179-187.
- Fujikawa, K., Chung, D.W., Hendrickson, L.E., Davie, E.W. (1986) Amino acid sequence of human factor XI, a blood coagulation factor with four tandem repeats that are highly homologous with plasma prekallikrein. Biochemistry. 25:2417-2424.
- Harris, J.R., Lippman, M.E., Veronesi, U.V., Willett, W. 1992 Breast Cancer N. Engl J Med 327:319-328, 390-398, 473-480.
- Kitamoto, Y., Veile, R.A., Donis-Keller, H. and Sadler, J.E. (1995) cDNA sequence and chromosomal localization of human enterokinase the proteolytic activator of trypsinogen. Biochemistry 34: 4562-4568.
- Lee, D.C., Fenton, S.E., Berkowitz, E.A. and Hissong, M.A. 1995 Transforming Growth Factor α : Expression, Regulation, and Biological Activities. Pharm. Rev. 47:51-85.
- Leytus, S.P., Loeb, K.R., Hagen, F.S., Kurachi, K., Davie, E.W. (1988) A novel trypsin-like serine protease (hepsin) with a putative transmembrane domain expressed by human liver and hepatoma cells. Biochemistry. 27:1067-1074.
- Miyazawa, K., Shimomura, T., Kitamura, A., Kondo, J., Morimoto, Y. and Kitamura, N. (1993) Molecular cloning and sequence analysis of the cDNA for a human serine protease responsible for activation of hepatocyte growth factor. Structural similarity of the protease precursor to blood coagulation factor XII. J. Biol. Chem. 268:10024-10028.
- Sakanari, J.A., Staunton, C.E., Eakin, A.E., Craik, C.S. and McKerrow, J.H. (1989) Serine Proteases from Nematode and Protozoan Parasites: Isolation of Sequence Homologs Using Generic Molecular Probes. Proc. Natl. Acad. Sci. USA. 86:4863-4867.
- Sambrook, J., Fritsch, E.F., Maniatis, T. Molecular Cloning: A Laboratory Manual, 2nd Edition, Cold Spring Harbor Laboratory Press, New York, 1989.
- Tamir, S., Kadner, S.S., Katz, J. and Finlay, T.H. 1990 Regulation of Antitrypsin and Antichymotrypsin Synthesis by MCF-7 Breast Cancer Cell Lines. Endocrinology 127:1319-1328.
- Yavelow J, Tuccillo A, Kadner SS, Katz J, Finlay TH. 1997 α_1 -Antitrypsin Blocks Release of TGF α from MCF-7 Human Breast Cancer Cells. J. Clin. Endo. Metabol 82:745-752.
- Yu, J.X., Chao, L. and Chao, J. 1994 Prostatin is a novel human serine proteinase from seminal fluid. Purification, tissue distribution, and localization in the prostate gland. J. Biol. Chem. 269:18843-18848.

Yu, J.X., Chao, L. and Chao, J. 1995 Molecular Cloning, tissue-specific expression and cellular localization of human prostatic mRNA. J. Biol. Chem 270:13483-13489.